

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 January 2004 (15.01.2004)

PCT

(10) International Publication Number
WO 2004/004693 A1

- (51) International Patent Classification⁷: **A61K 9/14**,
9/20, 9/24, 9/48 Roman, V.; 245 Kelton Street, Apartment 42, Allston, MA
02134 (US).
- (21) International Application Number: PCT/US2003/021095 (74) Agents: PABST, Patrea, L. et al.; Holland & Knight LLP,
One Atlantic Center, Suite 2000, 1201 West Peachtree
Street, N.E., Atlanta, GA 30309-3400 (US).
- (22) International Filing Date: 7 July 2003 (07.07.2003) (81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD,
SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG,
UZ, VC, VN, YU, ZA, ZM, ZW.
- (25) Filing Language: English (84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (26) Publication Language: English
- (30) Priority Data:
60/393,876 5 July 2002 (05.07.2002) US
60/436,523 23 December 2002 (23.12.2002) US
60/443,226 28 January 2003 (28.01.2003) US
60/463,514 15 April 2003 (15.04.2003) US
60/463,518 15 April 2003 (15.04.2003) US
- (71) Applicant: COLLEGIUM PHARMACEUTICAL
[US/US]; 400 Highland Corporate Drive, Cumberland, RI
02864 (US).
- (72) Inventors: HIRSH, Jane; 15 Peirce Road, Wellesley,
MA 02581 (US). KLIBANOV, Alexander, M.; 45 The
Ledges Road, Newton, MA 02460 (US). SWAGER,
Timothy, M.; 18 Copley Street, Newton, MA 02458
(US). BUCHWALD, Stephen, L.; 25 Lancaster Road,
Newton, MA 02458 (US). LO, Whe Yong; 65 Bayberry
Road, Canton, MA 02021 (US). FLEMING, Alison, B.;
8 Atlantic Street, Marshfield, MA 02050 (US). RARIY,
- Published:
— with international search report
— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments
- For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 2004/004693 A1

(54) Title: ABUSE-DETERRENT PHARMACEUTICAL COMPOSITIONS OF OPIOIDS AND OTHER DRUGS

(57) Abstract: An abuse-deterrent pharmaceutical composition has been developed to reduce the likelihood of improper administration of drugs, especially drugs such as opioids. In the preferred embodiment, a drug is modified to increase its lipophilicity. In preferred embodiments the modified drug is homogeneously dispersed within microparticles composed of a material that is either slowly soluble or not soluble in water. In some embodiments the drug containing microparticles or drug particles are coated with one or more coating layers, where at least one coating is water insoluble and preferably organic solvent insoluble, but enzymatically degradable by enzymes present in the human gastrointestinal tract. The abuse-deterrent composition retards the release of drug, even if the physical integrity of the formulation is compromised (for example, by chopping with a blade or crushing) and the resulting material is placed in water, snorted, or swallowed. However, when administered as directed, the drug is slowly released from the composition as the composition is broken down or dissolved gradually within the GI tract by a combination of enzymatic degradation, surfactant action of bile acids, and mechanical erosion.

ABUSE-DETERRENT PHARMACEUTICAL COMPOSITIONS OF OPIOIDS AND OTHER DRUGS

Background of the Invention

The present invention is generally in the field of pharmaceutical
5 compositions, and specifically relates to compositions that are designed to
reduce the potential for improper administration of drugs that are subject to
abuse.

This application claims priority to U.S.S.N.60/393,876 filed July 5,
2002 entitled "Abuse-Resistant Formulations of Oxycontin and Other Drugs"
10 by Alexander M. Klibanov, Stephen L. Buchwald, Timothy M. Swager, and
Whe-Yong Lo; U.S.S.N. 60/436,523 filed December 23, 2002 by Alison B.
Fleming, Roman V. Rariy, Alexander M. Klibanov, Whe-Yong Lo, and Jane
Hirsh; U.S.S.N. 60/443,226 filed January 28, 2003 by Jane Hirsh, Alison B.
Fleming, Alexander M. Klibanov, and Whe-Yong Lo; U.S.S.N. 60/463,514
15 filed April 15, 2003 by Jane C. Hirsh, Alison B. Fleming, Roman V. Rariy,
Stephen L. Buchwald, and Timothy M. Swager; and U.S.S.N. 60/463,518
filed April 15, 2003 by Jane C. Hirsh, Alison B. Fleming and Roman V.
Rariy.

Oxycodone, morphine, and other opiod analgesics are successful and
20 therapeutically useful medications, e.g., as pain killers, when administered
orally. Unfortunately, they also pose a severe threat for willful abuse due to
their ability to alter mood and/or cause a sense of euphoria. Currently
available sustained release formulations of such drugs, which contain a
relatively large amount of drug meant to be released from the formulation
25 over an extended time period, are particularly attractive to abusers since the
sustained release action can be destroyed by crushing or grinding the
formulation. The resulting material (ie, the crushed formulation) can no
longer control the release of drug. Depending on the drug, abusers can then
(1) snort the material, (2) swallow the material or (3) dissolve the material in
30 water and subsequently inject it intravenously. The dose of drug contained
in the formulation is thus absorbed immediately through the nasal or GI
mucosa (for snorting or swallowing, respectively) or is administered in a

bolus to the systemic circulation (for IV injection). These abuse methods result in the rapid bioavailability of relatively high doses of drug, giving the abuser a "high". Since relatively simple methods (crushing, grinding, chewing and/or dissolution in water) can be used to transform such formulations into an abusable form, they provide virtually no deterrent to a potential abuser.

For example, the FDA recently strengthened the warnings and precautions sections in the labeling of OxyContin® (oxycodone HCl controlled-release) Tablets, a narcotic drug approved for the treatment of moderate to severe pain, because of continuing reports of abuse and diversion. OxyContin® contains oxycodone HCl (available in 10, 20, 40 and 80 mg strengths), an opioid agonist with an addiction potential similar to that of morphine. Opioid agonists are substances that act by attaching to specific proteins called opioid receptors, which are found in the brain, spinal cord, and gastrointestinal tract. When these drugs attach to certain opioid receptors in the brain and spinal cord they can effectively block the transmission of pain messages to the brain. OxyContin® is supplied in a controlled-release dosage form and is intended to provide up to 12 hours of relief from moderate to severe pain. The warning specifically states that the tablet must be taken whole and only by mouth. When the tablet is chewed or crushed and its contents are swallowed, snorted into the nostrils or dissolved and subsequently injected intravenously, the controlled release mechanism is destroyed and a potentially lethal dose of oxycodone becomes bioavailable.

In recent years, there have been numerous reports of Oxycodone diversion and abuse in several states. For example, DEA's Office of Diversion Control reported 700 OxyContin® thefts in the US between January 2000 and June 2001. Some of these reported cases have been associated with serious consequences including death.

Oxycodone is a controlled substance in Schedule II of the Controlled Substances Act (CSA), which is administered by the Drug Enforcement Administration (DEA). Despite the fact that Schedule II provides the maximum amount of control possible under the CSA for approved drug

products, in practice it is difficult for law enforcement agencies to control the diversion or misuse of legitimate prescriptions. Although abuse, misuse, and diversion are potential problems for all opioids, including Oxycodone, opioids are a very important part of the medical armamentarium for the management of pain when used appropriately under the careful supervision of a physician.

Currently available formulations for such drugs are designed for oral administration but do not include mechanisms to prevent or retard improper methods of administration such as chewing, injection and snorting. This represents a serious problem given the large number of legitimate prescriptions written in the US; for example, the medical use of opioids within the US increased 400% from 1996 to 2000. The problems with abuse are significant and longstanding, and efforts to design new abuse-resistant or abuse-deterrent formulations have been largely unsuccessful. U.S. Patent numbers 3,980,766, 4,070,494 and 6,309,668 describe formulations designed to prevent the injection of compositions meant for oral administration. U.S. 3,980,766 describes the incorporation of an ingestible solid which causes a rapid increase in viscosity upon concentration of an aqueous solution thereof. U.S. 4,070,494 describes the incorporation of a non-toxic, water gelable material in an amount sufficient to render the drug resistant to aqueous extraction. U.S. 6,309,668 describes a tablet for oral administration containing two or more layers comprising one or more drugs and one or more gelling agents within separate layers of the tablet. The resulting tablet forms a gel when combined with the volume of water necessary to dissolve the drug; this formulation thus reduces the extractability of the drug from the tablet. It should be noted that although these compositions preclude abuse by injections, this approach fails to prevent abuse by crushing and swallowing or snorting the formulation, which are commonly reported methods of abuse associated with OxyContin®.

U.S. Patent numbers 3,773,955 and 3,966,940 describe formulations containing a combination of opioid agonists and antagonists, in which the antagonist does not block the therapeutic effect when the admixture is

administered orally, but which does not produce analgesia, euphoria or physical dependence when administered parenterally by an abuser. U.S. patent number 4,457,933 describes a method for decreasing both the oral and parenteral abuse potential of strong analgetic agents by combining an
5 analgesic dose of the analgetic agent with an antagonist in specific, relatively narrow ratios. U.S. Patent numbers 6,277,384, 6,375,957 and 6,475,494 describe oral dosage forms including a combination of an orally active opioid agonist and an orally active opioid antagonist in a ratio that, when delivered orally, is analgesically effective but that is aversive in a physically dependent
10 subject. While such a formulation may be successful in deterring abuse, it also has the potential to produce adverse effects in legitimate patients.

It is therefore an object of the present invention to provide a pharmaceutical composition that significantly reduces the potential for improper administration or use of drugs but which, when administered as
15 directed, is capable of delivering a therapeutically effective dose.

Summary of the Invention

An abuse-deterrent pharmaceutical composition and the method of making and using the composition have been developed. The composition can be used to reduce the likelihood of improper administration of drugs,
20 especially drugs such as oxycodone. The technology is useful for a number of other drugs where sustained release oral delivery is desired, and there is the potential for abuse if the drug dose is made immediately available for nasal, IV or oral administration. In the preferred embodiment, the drug is chemically modified to increase its lipophilicity. In other embodiments, the
25 formulation contains lipophilic or water-insoluble materials or is made using a process which increases the lipophilicity and/or water-insolubility of the composition.

The abuse-deterrent composition retards the release of drug, even if the physical integrity of the dosage form is compromised (for example, by
30 chopping with a blade or crushing) and the resulting material is placed in water, snorted, or swallowed. The composition thus provides a deterrent to common methods of improper administration including IV injection of drug

dissolved in water and nasal or oral administration of the crushed formulation since drug will not be immediately released from the formulation. However, when administered as directed, the drug is slowly released (typically over a period of 4-18 hours) from the composition as the composition is broken down or dissolved gradually within the GI tract by a combination of enzymatic degradation, surfactant action of bile acids, and mechanical erosion.

In some embodiments, the individual drug-containing microparticles or drug particles are coated with one or more independent coating layers. At least one of the coating materials is water-insoluble and preferably organic solvent-insoluble, but enzymatically degradable. The components of the resulting coated microparticles are not mutually soluble in water, organic solvents, or any combination thereof, so that *in vitro* degradation of the formulation will require more than one step. Hence the drug is not easily extractable from such a formulation by conventional chemical means. In contrast, when administered to the gastrointestinal tract via swallowing, the drug gradually will be released from the coated microparticles as a consequence of enzymatic degradation, surfactant action of bile acids and mechanical erosion within the GI tract.

The pharmaceutical composition, when administered orally, results in a desired drug release profile. Such a release profile provides a therapeutic effect for an extended period of time, typically from 6 to 24 hours. Additional compositions are provided which achieve a small immediate dose that precedes the sustained release of drug. The compositions disclosed herein may optionally comprise a drug having no appreciable abuse potential.

Brief Description of the Drawing

Figure 1A is a structural schematic of zinc bis-oxycodone. Figure 1B is a structural schematic of zinc oxycodone stearate. Figure 1C is a structural schematic of zinc oxycodone di-stearate.

Detailed Description of the Invention

Disclosed herein are an abuse-deterrent pharmaceutical composition and the method of making and using the composition.

I. Compositions

5 As used herein, "composition" refers to the drug dosage unit for administration to a patient. This may also be used in reference solely to the active ingredient, or to the formulation containing the active ingredient.

The currently available sustained release dosage forms containing narcotic analgesics and other drugs are subject to misuse, in part, because
10 mechanical destruction of the dosage form exposes the encapsulated drug and allows for immediate dissolution of the drug into aqueous media. Two properties of the dosage form that contribute to this outcome are (1) the ease with which drug is exposed to the extraction media and (2) the high water solubility of the drug salt form.

15 In the composition disclosed herein, one or both of these properties are altered in order to achieve an abuse-deterrent composition. Specifically, in the preferred embodiment, the drug is modified to increase its lipophilicity and, in additional preferred embodiments, is then homogeneously dispersed within a material that is either slowly soluble or not soluble in water and
20 subsequently formulated into microparticles. The drug may be present in the form of discrete particles or may be partially or fully dispersed in the carrier material on a molecular level.

The terms "abuse-deterrent composition" or "abuse-deterrent formulation" are used interchangeably herein to refer to compositions that
25 reduce the potential for improper administration of drugs but that deliver a therapeutically effective dose when administered as directed. Improper administration includes tampering with the dosage form and/or administering the drug by any route other than instructed. For example, for a tablet or capsule, methods of tampering with the dosage form may include, but are not
30 limited to, breaking, crushing, grinding, chewing and/or dissolving the tablet or the contents of the capsule. For oral administration, improper

administration includes administering the drug by any route other than via swallowing.

The abuse deterrent composition preferably comprises a drug modified to increase its lipophilicity. In other preferred embodiments, the drug is homogeneously dispersed within microparticles composed of a material that is either slowly soluble in water or water insoluble. The compositions slow the release of drug if the dosage form is chopped or crushed and the resulting material is placed in water, snorted, or swallowed since most of the drug will remain associated with or entrapped within portions of the core material of the microparticles. In some embodiments the drug containing microparticles or individual drug particles are coated with one or more coating layers, where at least one coating is water insoluble and preferably organic solvent insoluble, but enzymatically degradable. The components of the resulting coated microparticles are not mutually soluble in water, organic solvents, or any combination thereof, such that no one solvent or enzyme solution is capable of dissolving the formulation in its entirety *in vitro*. It follows that extraction of the drug from the formulation cannot be carried out in one step. However, when administered as directed, the drug is slowly released from the formulation since it is eroded within the environment of the gastrointestinal tract.

A. Drugs to be formulated

There are many drugs that it is desirable to deliver using the compositions described herein. The Controlled Substances Act (CSA), Title II of the Comprehensive Drug Abuse Prevention and Control Act of 1970, places all substances that are regulated under existing federal law into one of five schedules based upon the substance's medicinal value, harmfulness, and potential for abuse or addiction. Drugs that are preferred include those classified as Schedule II, III, IV and V drugs. Drugs that are most preferable include those, like oxycodone, that are currently formulated as sustained or controlled release compositions, where drug release is intended to occur over a prolonged period of time through the gastrointestinal tract, and immediate or burst release, for example, by inhalation or injection, is undesirable. As

used herein, drugs prone to abuse refer to controlled substance specified as schedule II, III, IV and V drugs.

The terms "drug", "active agent", and "pharmacologically active agent" are used interchangeably herein to refer to a chemical compound that induces a desired pharmacological, physiological effect. The terms also encompass pharmaceutically acceptable derivatives of those active agents specifically mentioned herein, including, but not limited to, salts, solvates, hydrates, complexes with one or more molecules, prodrugs, active metabolites, analogs, and the like. When the terms "active agent", "pharmacologically active agent" and "drug" are used, or when a particular drug, such as oxycodone, is identified, it is to be understood as including the active agent per se as well as pharmaceutically acceptable salts, solvates, hydrates, complexes with one or more molecules, prodrugs, active metabolites, and analogs.

Examples of preferred drugs include, 1-phenylcyclohexylamine, 1-piperidinocyclohexanecarbonitrile, alfentanil, alphacetylmethadol, alphaprodine, alprazolam, amobarbital, amphetamine, anileridine, apomorphine, aprobarbital, barbital, barbituric acid derivative, bemidone, benzoylecgonine, benzphetamine, betacetylmethadol, betaprodine, bezitramide, bromazepam, buprenorphine, butabarbital, butalbital, butorphanol, camazepam, cathine, chloral, chlordiazepoxide, clobazam, clonazepam, clorazepate, clotiazepam, cloxazolam, cocaine, codeine, chlorphentermine, delorazepam, dexfenfluramine, dextromoramide, dextropropoxyphen, dezocine, diazepam, diethylpropion, difenoxin, dihydrocodeine, dihydromorphine, dioxaphentyl butyrate, dipanone, diphenoxylate, diprenorphine, ecgonine, enadoline, eptazocine, estazolam, ethoheptazine, ethyl loflazepate, ethylmorphine, etorphine, fempropionex, fencamfamin, fenfluramine, fentanyl, fludiazepam, flunitrazepam, flurazepam, glutethimide, halazepam, haloxazolam, hexalgon, hydrocodone, hydromorphone, isomethadone, hydrocodone, ketamine, ketazolam, ketobemidone, levanone, levoalphacetylmethadol, levomethadone, levomethadyl acetate, levomethorphan, levorphanol, lofentanil, loperamide,

loprazolam, lorazepam, lormetazepam, lysergic acid, lysergic acid amide,
mazindol, medazepam, mefenorex, meperidine, meptazinol, metazocine,
methadone, methamphetamine, methohexital, methotrimeprazine,
methyldihydromorphinone, methylphenidate, methylphenobarbital, metopon,
5 morphine, nabilone, nalbuphine, nalbupine, nalorphine, narceine, nefopam,
nicomorphine, nimetazepam, nitrazepam, nordiazepam, normethadone,
normorphine, oxazepam, oxazolam, oxycodone, oxymorphone, pentazocine,
pentobarbital, phenadoxone, phenazocine, phencyclidine, phendimetrazine,
phenmetrazine, pheneridine, piminodine, prodilidine, properidine,
10 propoxyphene, racemethorphan, racemorphan, racemoramide, remifentanil,
secobarbital, sufentanil, talbutal, thebaine, thiamylal, thiopental, tramadol,
trimeperidine, and vinbarbital.

In addition to the compounds above, the following scheduled drugs
may be incorporated into the composition: allobarbitone, alprazolam,
15 amylobarbitone, aprobarbital, barbital, barbitone, benzphetamine,
brallobarbital, bromazepam, brotizolam, buspirone, butalbital, butobarbitone,
butorphanol, camazepam, captodiame, carbromal, carfentanil, carpipramine,
cathine, chloral, chloral betaine, chloral hydrate, chloralose,
chlordiazepoxide, chlorhexadol, chlormethiazole edisylate, chlormezanone,
20 cinolazepam, clobazam, potassium clorazepate, clotiazepam, cloxazolam,
cyclobarbitone, delorazepam, dexfenfluramine, diazepam, diethylpropion,
difebarbamate, difenoxin, enciprazine, estazolam, ethyl loflazepate, etizolam,
febarbamate, fencamfamin, fenfluramine, fenproporex, fluanisone,
fludiazepam, flunitraam, flunitrazepam, flurazepam, flutoprazepam,
25 gepirone, glutethimide, halazepam, haloxazolam, hexobarbitone, ibomal,
ipsapirone, ketazolam, loprazolam mesylate, lorazepam, lormetazepam,
mazindol, mebutamate, medazepam, mefenorex, mephobarbital,
meprobamate, metaclozepam, methaqualone, methohexital, methylpentynol,
methylphenobarbital, midazolam, milazolam, morphine, nimetazepam,
30 nitrazepam, nordiazepam, oxazepam, oxazolam, paraldehyde, pemoline,
pentabarbitone, pentazocine, pentobarbital, phencyclidine, phenobarbital,
phendimetrazine, phenmetrazine, phenprobamate, phentermine,

phenylacetone, pinazepam, pipradol, prazepam, proxibarbal, quazepam, quinalbaritone, secobarbital, secbutobarbitone, sibutramine, temazepam, tetrazepam, triazolam, triclofos, zalepan, zaleplon, zolazepam, zolpidem, and zopiclone.

- 5 Certain compounds described herein may exist in particular geometric or stereoisomeric forms. The composition disclosed herein contemplates all such compounds, including *cis*- and *trans*-isomers, *R*- and *S*-enantiomers, diastereomers, (D)-isomers, (L)-isomers, the racemic mixtures thereof, compounds of different spacial conformations, and other mixtures thereof, as
10 falling within the scope of the invention. Additional asymmetric carbon atoms may be present in a substituent such as an alkyl group. All such isomers, as well as mixtures thereof, are intended to be included in this invention.

- As used herein, "pharmaceutically acceptable salts" refer to
15 derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically
20 acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts
25 prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, tolunesulfonic, methanesulfonic, ethane disulfonic, oxalic, and isethionic.

- 30 The pharmaceutically acceptable salts of the compounds can be synthesized from the parent compound, which contains a basic or acidic moiety, by conventional chemical methods. Generally, such salts can be

prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, non-aqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred.

- 5 Lists of suitable salts are found in Remington's Pharmaceutical Sciences, 20th ed., Lippincott Williams & Wilkins, Baltimore, MD, 2000, p. 704, the disclosure of which is hereby incorporated by reference.

Optionally, the composition described herein can further include a drug having no appreciable abuse potential.

10 **B. Drug solubility modification**

- In preferred embodiments, the solubility characteristics of a drug are altered prior to incorporation into the formulation. Modification of the drug to produce a more lipophilic derivative serves to reduce the water solubility of the drug and thus reduces the aqueous extractability. Furthermore, if the drug is made more lipophilic, it can be solubilized in a molten fatty substance or wax like mixture, rather than physically dispersed in a particulate form. Solubilization of drug enhances the abuse-deterrent properties of microparticles formulated from the mixture as it is difficult to extract drug from an intimately dispersed composition.

- 20 The terms "lipophilic derivative" and "lipophilic drug derivative", as used herein, refer to derivatives of the drug that are less soluble in water than the most soluble salt of the drug. The most soluble salt being selected from either drug alkaline metal salts (for acidic drugs) or salts of the drug with inorganic acids (for basic drugs). The examples of the latter include but are not limited to hydrohalates, sulfates, and nitrates.

Some of the methods that can be used to alter the drug's lipophilicity are outlined below. It is understood that two or more approaches can be combined to achieve a desired solubility profile.

Methods for increasing lipophilicity

- 30 In one embodiment, drug is made more lipophilic by eliminating or reducing the overall charge of the drug molecule. For example, for a basic drug, a water soluble salt (such as hydrochloride, sulfate, or maleate) can be

converted to a free base using techniques known in the art. Correspondingly, in the case of an acidic drug, a water soluble salt (such sodium, potassium, or the like) can be converted to a free acid.

In another embodiment, the drug's lipophilicity is increased by forming a salt between a drug molecule and a charged lipophilic compound. In this case the lipophilicity of the resulting salt can be manipulated by varying the lipophilicity of the counter-ion. In general lipophilic acids or amines with chain lengths between C₅-C₃₀ are lipophilic counter-ion candidates. Some specific examples include, but are not limited to, linoleic acid, octanoic acid, lauric acid, stearic acid, palmitic acid, oleic acid, octyl amine, lauryl amine, stearyl amine, palmityl amine, linoleyl amine, and oleyl amine. Other salts which may increase lipophilicity and, hence, lipid solubility relative to the parent drug compound include, but are not limited to, pectinate, tannate, phytate, salicylate, saccharinate, acesulfamate, gallate, and terephthalate salts.

In yet another embodiment the lipophilicity of the drug is increased by forming a stable complex between a drug molecule (either charged or uncharged) and a metal cation such as zinc, magnesium, calcium, bismuth or the like. This complex may consist of one or more drug molecules, one or more metal cations, and, optionally, one or more lipophilic charged species. The aforementioned charged lipophilic species are incorporated into the complex if necessary to bring the charge of the final complex to zero and increase its overall lipophilicity. In general lipophilic acids or amines with chain lengths between C₅-C₃₀ are lipophilic counter-ion candidates. Examples of such complexes for a narcotic drug oxycodone are given in Figure 1; a lipophilic drug complex may be composed of one or two oxycodone molecules, one Zn²⁺ cation, and one or two stearate anions. It is understood by one skilled in the art that various metal cations as well as lipophilic counter-ions can be used to form complexes with an analogous structure, for example, oxymorphone.

In still a further embodiment, drug lipophilicity is increased via complexation with poorly water-soluble cyclodextrin. For example,

ethylated beta-cyclodextrin has been shown to decrease aqueous solubility of complexed drug molecules.

In another embodiment, a drug is covalently modified to increase its lipophilicity. For example, a lipophilic compound can be covalently attached to a drug molecule via an ester or amide linkage. Such drug derivatives are
5 cleaved *in vivo*, thus releasing the parent compound.

C. Drug Containing Microparticles

In preferred embodiments, drugs are formulated with a carrier material to form microparticles. As used herein, the term "microparticle"
10 refers to a composition comprising a drug dispersed within a carrier material and "coated microparticle" refers to a composition comprising a drug containing microparticle or a drug particle coated with one or more coating layers of material. Microparticles and coated microparticles have a size range of 10 to 3000 microns in diameter.

15 Within microparticles, drug is preferably homogeneously dispersed in the form of fine particles within the carrier material. More preferably, drug is partially solubilized in molten carrier material or partially dissolved with the carrier material in a mutual solvent during the formulation of the microparticles. Most preferably, drug is completely solubilized in the molten
20 carrier material or completely dissolved with the carrier material in a co-solvent during the formulation of the microparticles. This is accomplished through the selection of materials and the manner in which they are processed.

Carrier materials appropriate for the fabrication of drug containing
25 microparticles are either slowly soluble in water or insoluble in water, but capable of degrading within the GI tract by means including enzymatic degradation, surfactant action of bile acids and mechanical erosion. As used herein, the term "slowly soluble in water" refers to materials that are not dissolved in water within a period of 30 minutes. Preferred examples include
30 fats, fatty substances, waxes, wax-like substances and mixtures thereof. Suitable fats and fatty substances include fatty alcohols (such as lauryl, myristyl stearyl, cetyl or cetostearyl alcohol), fatty acids and derivatives,

including but not limited to fatty acid esters, fatty acid glycerides (mono-, di- and tri-glycerides), and hydrogenated fats. Specific examples include, but are not limited to hydrogenated vegetable oil, hydrogenated cottonseed oil, hydrogenated castor oil, hydrogenated oils available under the trade name
5 Sterotex®, stearic acid, cocoa butter, and stearyl alcohol. Suitable waxes and wax-like materials include natural or synthetic waxes, hydrocarbons, and normal waxes. Specific examples of waxes include beeswax, glycowax, castor wax, carnauba wax, paraffins and candelilla wax. As used herein, a wax-like material is defined as any material which is normally solid at room
10 temperature and has a melting point of from about 30 to 300°C.

In some cases, it may be desirable to alter the rate of water penetration into the hydrophobic drug containing microparticles. To this end, rate-controlling (wicking) agents may be formulated along with the fats or waxes listed above. Examples of rate-controlling materials include certain
15 starch derivatives (eg, waxy maltodextrin and drum dried corn starch), cellulose derivatives (eg, hydroxypropylmethylcellulose, hydroxypropylcellulose, methylcellulose, and carboxymethylcellulose), alginic acid, lactose and talc. Additionally, a pharmaceutically acceptable surfactant (for example, lecithin) may be added to facilitate the degradation
20 of such microparticles.

Proteins which are water insoluble, such as zein, are preferred carrier materials for the formation of drug containing microparticles. Additionally, proteins, polysaccharides and combinations thereof which are water soluble can be formulated with drug into microparticles and subsequently cross-
25 linked to form an insoluble network. For example, cyclodextrins can be complexed with individual drug molecules and subsequently cross-linked.

Certain polymers may also be used as carrier materials in the formulation of drug containing microparticles. Suitable polymers include ethylcellulose and other natural or synthetic cellulose derivatives. Polymers
30 which are slowly soluble and form a gel in an aqueous environment, such as hydroxypropyl methylcellulose or polyethylene oxide may also be suitable as carrier materials for drug containing microparticles.

Encapsulation or incorporation of drug into carrier materials to produce drug containing microparticles can be achieved through known pharmaceutical formulation techniques. To create a composition that protects drug from exposure upon mechanical disruption (eg, grinding, chewing, or chopping), the drug is intimately dispersed within the carrier material. In the case of formulation in fats, waxes or wax-like materials, the carrier material is heated above its melting temperature and the drug is added to form a mixture comprising drug particles suspended in the carrier material, drug dissolved in the carrier material, or a mixture thereof.

10 Microparticles can be subsequently formulated through several methods including, but not limited to, the processes of congealing, extrusion, spray chilling or aqueous dispersion. In a preferred process, wax is heated above its melting temperature, drug is added, and the molten wax-drug mixture is congealed under constant stirring as the mixture cools. Alternatively, the

15 molten wax-drug mixture can be extruded and spheronized to form pellets or beads. Detailed descriptions of these processes can be found in "Remington-The science and practice of pharmacy", 20th Edition, Jennaro et. Al., (Phila, Lippencott, Williams, and Wilkens, 2000).

For some carrier materials it may be desirable to use a solvent evaporation technique to produce drug containing microparticles. In this case drug and carrier material are co-dissolved in a mutual solvent and microparticles can subsequently be produced by several techniques including, but not limited to, forming an emulsion in water or other appropriate media, spray drying or by evaporating off the solvent from the

25 bulk solution and milling the resulting material.

In addition to modification of the drug itself, processing conditions can be used to influence the dispersion of the drug within water-insoluble or slowly water soluble material. For example, in the case where the water insoluble or slowly soluble material is melted and drug is fully or partially

30 dissolved under stirring conditions, the temperature, agitation rate and time of processing will influence the degree of dissolution achieved. More specifically, a more homogenous dispersion may be achieved with a higher

temperature, faster stirring rate and longer processing time. Ultrasound can also be applied to the molten mixture to increase the degree of dispersion and/or the rate of dissolution of the drug.

In some embodiments, drug in a particulate form is homogeneously dispersed in a water-insoluble or slowly water soluble material. To minimize the size of the drug particles within the composition, the drug powder itself may be milled to generate fine particles prior to formulation. The process of jet milling, known in the pharmaceutical art, can be used for this purpose. In some embodiments drug in a particulate form is homogeneously dispersed in a wax or wax like substance by heating the wax or wax like substance above its melting point and adding the drug particles while stirring the mixture. In this case a pharmaceutically acceptable surfactant may be added to the mixture to facilitate the dispersion of the drug particles.

D. Coated Drug Containing Microparticles

In some embodiments, drug containing microparticles or drug particles are encapsulated within at least one water-insoluble enzymatically degradable material. In some instances the substrates of digestive enzymes are naturally water-insoluble and can be utilized in the formulation without further processing. Solid esters of fatty acids, which are hydrolyzed by lipases, can be spray coated onto microparticles or drug particles. Zein is an example of a naturally water-insoluble protein. It can be coated onto drug containing microparticles or drug particles by spray coating or by wet granulation techniques. In addition to naturally water-insoluble materials, some substrates of digestive enzymes can be treated with cross-linking procedures, resulting in the formation of non-soluble networks. Many methods of cross-linking proteins, initiated by both chemical and physical means, have been reported. One of the most common methods to obtain cross-linking is the use of chemical cross-linking agents. Examples of chemical cross-linking agents include aldehydes (gluteraldehyde and formaldehyde), epoxy compounds, carbodiimides, and genipin. In addition to these cross-linking agents, oxidized and native sugars have been used to cross-link gelatin (Cortesi, R., et al., Biomaterials 19 (1998) 1641-1649).

Cross-linking can also be accomplished using enzymatic means; for example, transglutaminase has been approved as a GRAS substance for cross-linking seafood products. Finally, cross-linking can be initiated by physical means such as thermal treatment, UV irradiation and gamma
5 irradiation.

To produce a coating layer of cross-linked protein surrounding drug containing microparticles or drug particles, a water soluble protein can be spray coated onto the microparticles and subsequently cross-linked by the one of the methods described above. Alternatively, drug containing
10 microparticles can be microencapsulated within protein by coacervation-phase separation (for example, by the addition of salts) and subsequently cross-linked. Some suitable proteins for this purpose include gelatin, albumin, casein, and gluten.

Polysaccharides can also be cross-linked to form a water-insoluble
15 network. For many polysaccharides, this can be accomplished by reaction with calcium salts or multivalent cations which cross-link the main polymer chains. Pectin, alginate, dextran, amylose and guar gum are subject to cross-linking in the presence of multivalent cations. Complexes between oppositely charged polysaccharides can also be formed; pectin and chitosan,
20 for example, can be complexed via electrostatic interactions. Insoluble coatings can be formed on particles in this fashion. It should be noted that in many cases polysaccharides are broken down specifically by enzymes produced by bacteria within the colon.

In some cases a water-insoluble but enzymatically degradable coating
25 comprising both a protein and a polysaccharide can be produced if the components are oppositely charged polyelectrolytes. Under the proper temperature, pH, and concentrations, the two polymers can interact through their opposite electrical charges and form a water-insoluble complex. If a core particle is present at the time the complex phase separates, it will be
30 coated. For example, gelatin and gum arabic can be coated onto a core particle utilizing this process. Optionally, the complex can be made

irreversibly insoluble by subsequent cross-linking induced by chemical or physical means.

E. Dosage Forms

There are a number of drug compositions that meet the abuse
5 deterrent criteria outlined above. In one embodiment a drug is
homogeneously dispersed, in a fine particulate form, within a water-insoluble
or slowly water soluble material and the mixture is formulated into
microparticles. In another embodiment a drug is partially dissolved within a
10 water-insoluble or slowly water soluble material during the manufacturing
process, for example, by mixing at a temperature above the melting point of
the carrier material, and the mixture is formulated into microparticles. In yet
another embodiment a drug is fully dissolved within a water-insoluble or
slowly water soluble material during the manufacturing process, for example,
15 by mixing at a temperature above the melting point of the carrier material,
and the mixture is formulated into microparticles. In still a further
embodiment, the drug containing microparticles, where the drug is
homogeneously dispersed in a particulate form, or has been partially or fully
dissolved within the carrier material during the manufacturing process, are
coated with one or more coatings to form coated microparticles. In a further
20 embodiment, drug particles are coated directly with one or more coatings to
form coated microparticles.

The microparticles, coated microparticles, or a mixture thereof are
formed into a solid dosage form suitable for oral administration. For
example, microparticles or coated microparticles can be incorporated into
25 hard capsules, dispersed within a soft gelatin capsule, or combined with
appropriate excipients and tableted by compression.

In some embodiments, the compositions are coated with an enteric
coating. Enteric coatings known in the art are applied directly to the abuse-
deterrent microparticle or coated microparticle compositions or are applied to
30 the surface of a capsule or tablet comprising the abuse deterrent
microparticle and/or coated microparticle compositions. Enteric coatings
known in the art include, for example, acrylic polymers that are

commercially available under the trade name EUDRAGIT ®, cellulose acetate phthalate, hydroxypropylmethylcellulose phthalate, polyvinylacetate phthalate, shellac, hydroxypropylmethylcellulose succinate, cellulose acetate trimellitate or mixtures thereof.

5 Dosage forms can include one or more drugs. When the dosage form includes two or more drugs they can be Scheduled drugs or can be a combination of Scheduled and non-Scheduled drugs. The drugs can be incorporated into separate microparticle compositions where the Scheduled drugs are incorporated into abuse deterrent microparticle compositions and
10 the non-Scheduled drugs are incorporated into abuse deterrent microparticle compositions, sustained release compositions known in the art or immediate release compositions known in the art. The compositions comprising the different drugs are formulated into a single solid dosage form suitable for oral administration, for example, they can be incorporated into a gelatin
15 capsule, or combined with appropriate excipients and compressed into a tablet form. Examples of non-scheduled drugs that may be included in dosage forms described herein include, but are not limited to, aspirin, acetaminophen, non-steroidal anti-inflammatory drugs, cyclooxygenase II inhibitors, N-methyl-D-aspartate receptor antagonists, glycine receptor
20 antagonists, triptans, dextromethorphan, promethazine, fiorinal, guaifenesin, butalbital, and caffeine.

 An immediate release dose can be incorporated into the formulation in several ways. Immediate release microparticles can be made utilizing standard methodologies and formulated along with abuse-deterrent
25 microparticle and/or coated microparticle compositions in a suitable oral dosage form. Alternatively, a coating containing drug which is available for immediate release can be placed on a tablet comprising abuse-deterrent microparticle and/or coated microparticle compositions plus appropriate excipients. Additionally, an immediate dose of drug can be granulated or
30 blended with rapidly dissolving excipients and subsequently compressed (1) as one layer of bi-layer tablets in which the abuse-deterrent microparticle and/or coated microparticle compositions are compressed as the other layer,

or (2) as the outer layer of compression-coated tablets in which the abuse-deterrent microparticle and/or coated microparticle compositions are compressed as the inner core, or (3) into tablets in which abuse-deterrent microparticle and/or coated microparticle compositions are embedded.

5 In some embodiments, the immediate release portion of the dosage form comprises a lipophilic drug derivative. For example, salt derivatives or complexes that are insoluble at a neutral pH but dissociate, thereby releasing the parent compound, at an acidic pH are ideal for immediate release within the stomach. In the case of oxycodone some salts that may exhibit this
10 property include, but are not limited to, the tannate, phthalate, salicylate, gallate, pectinate, phytate, saccharinate, asesulfamate and terephthalate salts. Complexes of drug with one or more metal ions and, optionally, one or more lipophilic counter-ions (see, for example, Figure 1) may also be used for immediate drug release. Use of salts or complexes in the immediate release
15 portion of the dosage form reduces the abuse potential of the immediate release dose if the formulation is crushed and (1) snorted or (2) dissolved in water since these salts will be poorly soluble under these conditions. It is understood by the one of ordinary skill in the art that such salts or complexes may also be used to formulate an immediate release dosage form without a
20 sustained release portion.

 Additional mechanisms to reduce the potential for abuse can also be incorporated during the process of formulating tablets. For example, ingredients can be added to deter chewing or snorting of the final formulation. For example, an intensely bitter substance may deter chewing,
25 while an intensely spicy ingredient, such as capsaicin, may deter snorting. The addition of a colored dye, which would stain the skin and mucosal surface of the nose following snorting may also serve to reduce this practice.

 Optional excipients present in the oral dosage form comprising abuse deterrent microparticles or coated microparticles include, but are not limited
30 to diluents, binders, lubricants, disintegrants, colorants, plasticizers and the like. Diluents, also termed "fillers," are typically necessary to increase the bulk of a solid dosage form so that a practical size is provided for

compression of tablets. Examples of diluents include cellulose, dry starch, microcrystalline cellulose, dicalcium phosphate, calcium sulfate, sodium chloride confectioner's sugar, compressible sugar, dextrans, dextrin, dextrose, sucrose, mannitol, powdered cellulose, sorbitol, and lactose.

5 Binders are used to impart cohesive qualities powdered materials and can include materials such as starch, gelatin, sugars, natural and synthetic gums, polyethylene glycol, ethylcellulose, methylcellulose, hydroxypropylmethylcellulose, carboxymethylcellulose, waxes and polyvinyl pyrrolidone. Lubricants are used to facilitate tablet manufacture;

10 examples of lubricants include talc, magnesium stearate, calcium stearate, hydrogenated vegetable oils stearic acid, sodium stearyl fumarate, sodium benzoate, sodium acetate, leucine, sodium oleate, sodium lauryl sulfate, magnesium lauryl sulfate and polyethylene glycol. Disintegrants can be added to pharmaceutical formulations in order to facilitate "breakup" or

15 disintegration after administration. Materials used for this purpose include starches, clays, celluloses, alginates, gums, and cross-linked polymers. A plasticizer may be included in coating materials to alter their mechanical properties. Examples of plasticizers include benzyl benzoate, chlorobutanol, dibutyl sebacate, diethyl phthalate, glycerin, mineral oil, polyethylene glycol,

20 sorbitol, triacetin, triethyl citrate, glycerol, etc. In addition to the additives above, coloring and flavoring agents may also be incorporated into the composition.

Optionally, the composition disclosed herein comprises materials wherein a combination of the materials is not soluble in water, organic

25 solvent, or any combination thereof.

II. Methods of Administration

It is assumed that upon oral ingestion of the intact composition, drug is released as the formulation is gradually broken down or dissolved within the GI tract by a combination of enzymatic degradation, surfactant action of

30 bile acids, and mechanical erosion. This is a result of the unique ability of the human digestive system to efficiently break down or solubilize a variety of materials. The process within the GI tract that results in the digestion of

food and the absorption of nutrients is well known. Following mastication within the mouth, food passes into the stomach where it is mixed with digestive juices. This fluid contains the proteolytic enzyme pepsin which, following activation by the low pH within the stomach, begins the process of

5 cleaving ingested proteins into smaller peptide fragments. Food then enters the small intestine in the form of macromolecular aggregates, where it is digested into molecules near or in a form capable of being absorbed. This digestion is accomplished through the action of various enzymes which are produced in the pancreas and flow into the upper portion of the large

10 intestine, the duodenum. The enzymes synthesized in the pancreas include proteases, amylases and lipases; these enzymes are capable of breaking down proteins, starches and fats, respectively. The digestion of fats is further facilitated by the secretion of bile into the duodenum since bile salts, which contain both hydrophobic and hydrophilic portions, are capable of

15 emulsifying lipids into minute droplets in order to increase the surface area available for digestion by lipases. The material which remains following passage through the small intestine enters the large intestine. Bacteria capable of breaking down carbohydrates not digested in the small intestine (such as cellulose) are present in large numbers this region of the digestive

20 tract. Finally, in addition to microbial fermentation, the large intestine functions to absorb water and electrolytes and to form and store feces until they are excreted.

In addition to providing a deterrent to common methods of abuse/diversion, the formulation can provide a sustained release of drug over

25 an extended time period. This is a natural consequence of the fact that, in the present formulation, drug is slowly released from a predominantly water-insoluble, hydrophobic matrix following the degradation of the matrix. The barrier components are degraded, for example, by enzymes, the surfactant action of bile acids and mechanical erosion.

30 In some embodiments, an immediate release of drug is achieved within the stomach in order to provide rapid therapeutic onset.

The pharmaceutical drug composition is administered orally. The appropriate dosage formulations can be obtained by calculation of the pharmacokinetics of the formulation, then adjusting using routine techniques to yield the appropriate drug levels based on the approved dosage forms. Any
5 suitable amount of drug containing microparticles or coated microparticles can be included in the final formulation. The selection of a suitable amount of drug containing microparticles depends on the dosage desired and is readily determined by those skilled in the art.

In addition to oral administration, some embodiments may also be
10 administered by other routes, including, but not limited to, rectal and nasal administration. Some embodiments may also be suitable for formulation as oral liquids.

The present composition and method of making and using the composition will be further understood by reference to the following non-
15 limiting examples.

Example 1: Preparation of lipophilic oxycodone derivatives.

A. Oxycodone free base

The free base of oxycodone was prepared from its hydrochloride salt by the following method: Oxycodone hydrochloride was dissolved in water
20 and sodium carbonate was added in the amount required to neutralize hydrochloric acid. Methylene chloride was added in order to extract the formed oxycodone free base. The obtained organic layer was dried over sodium sulfate and methylene chloride was evaporated using rotary
evaporator. The obtained oxycodone free base was purified by
25 crystallization.

B. Zinc-bis-oxycodone

Zinc bis-oxycodone can be obtained in anhydrous media by reacting oxycodone free base with $\text{Zn}(\text{Et})_2$

C. Zinc oxycodone stearate

30 Zinc oxycodone stearate can be obtained in anhydrous media by reacting oxycodone free base with $\text{Zn}(\text{Et})(\text{C}_{18}\text{H}_{35}\text{O}_2)$

D. Zinc oxycodone di-stearate

.Zinc-oxycodone di-stearate can be obtained by co-melting
 $\text{Zn}(\text{C}_{18}\text{H}_{35}\text{O}_2)_2$ and oxycodone free base.

E. Oxycodone terephthalate

Oxycodone terephthalate is commercially available and can be used
5 without further processing

The structures of some representative oxycodone zinc complexes are
shown in Figures 1A, 1B and 1C.

Example 2: Preparation of drug containing microparticles

The free base, salts or complexes from Example 1 are added to
10 molten hydrogenated vegetable oil, mixed, extruded and spheronized to form
drug containing microparticles.

Example 3: Preparation of coated drug containing microparticles

The drug-containing particles from Example 2 are spray coated with
zein in a fluidized bed apparatus.

15 **Example 4: Preparation of capsules for oral administration**

The drug containing microparticles from Example 2 and/or the coated
microparticles from Example 3 are incorporated into standard gelatin
capsules.

20 **Example 5: Preparation of capsules for oral administration containing a
dose of drug for immediate release**

The drug containing microparticles from Example 2 and/or the coated
microparticles from Example 3 are combined with immediate release drug
particles, and incorporated into standard gelatin capsules.

25

We claim:

1. An orally administerable abuse-deterrent pharmaceutical composition comprising a therapeutically effective amount of a drug prone to abuse selected from the group of compositions consisting of
 - (a) a composition comprising a therapeutically effective amount of a lipophilic derivative of a drug prone to abuse, and
 - (b) a water-insoluble, preferably lipophilic, formulation comprising a therapeutically effective amount of a drug prone to abuse.
2. The composition of claim 1 comprising a therapeutically effective amount of a lipophilic derivative of a drug prone to abuse in one or more pharmaceutically acceptable excipients.
3. The composition of Claim 1, wherein the composition is a controlled-release pharmaceutical composition.
4. The composition of Claim 1, wherein the composition prevents the immediate release of a substantial portion of incorporated drug when the physical integrity of the composition is compromised and the resulting material is exposed to an aqueous medium.
5. The composition of Claim 4, wherein the portion of the drug released immediately is less than 80% of the total amount of drug incorporated into formulation.
6. The composition of Claim 1, wherein the composition prevents the immediate release of a substantial portion of incorporated drug when the physical integrity of the composition is compromised and the resulting material is exposed to a non-aqueous medium.
7. The composition of Claim 6, wherein the portion of the drug released immediately is less than 80% of the total amount of the drug incorporated into the composition.
8. The composition of Claim 1 wherein the drug prone to abuse is selected from the group consisting of 1-phenylcyclohexylamine, 1-piperidinocyclohexanecarbonitrile, alfentanil, alphacetylmethadol, alphaprodine, alprazolam, amobarbital, amphetamine, anileridine, apomorphine, aprobarbital, barbital, barbituric acid derivative, bemidone,

benzoylecgonine, benzphetamine, betacetylmethadol, betaprodine, bezitramide, bromazepam, buprenorphine, butabarbital, butalbital, butorphanol, camazepam, cathine, chloral, chlordiazepoxide, clobazam, clonazepam, clorazepate, clotiazepam, cloxazolam, cocaine, codeine, chlorphentermine, delorazepam, dexfenfluramine, dextromoramide, dextropropoxyphen, dezocine, diazepam, diethylpropion, difenoxin, dihydrocodeine, dihydromorphine, dioxaphentyl butyrate, dipanone, diphenoxylate, diprenorphine, ecgonine, enadoline, eptazocine, estazolam, ethoheptazine, ethyl loflazepate, ethylmorphine, etorphine, femproporex, fencamfamin, fenfluramine, fentanyl, fludiazepam, flunitrazepam, flurazepam, glutethimide, halazepam, haloxazolam, hexalgon, hydrocodone, hydromorphone, isomethadone, hydrocodone, ketamine, ketazolam, ketobemidone, levanone, levoalphacetylmethadol, levomethadone, levomethadyl acetate, levomethorphan, levorphanol, lofentanil, loperamide, loprazolam, lorazepam, lormetazepam, lysergic acid, lysergic acid amide, mazindol, medazepam, mefenorex, meperidine, meptazinol, metazocine, methadone, methamphetamine, methohexital, methotrimeprazine, methyl dihydromorphinone, methylphenidate, methylphenobarbital, metopon, morphine, nabilone, nalbuphine, nalbupine, nalorphine, narceine, nefopam, nicomorphine, nimetazepam, nitrazepam, nordiazepam, normethadone, normorphine, oxazepam, oxazolam, oxycodone, oxymorphone, pentazocine, pentobarbital, phenadoxone, phenazocine, phencyclidine, phendimetrazine, phenmetrazine, pheneridine, piminodine, prodilidine, properidine, propoxyphene, racemethorphan, racemorphan, racemoramide, remifentanyl, secobarbital, sufentanil, talbutal, thebaine, thiamylal, thiopental, tramadol, trimeperidine, vinbarbital, allobarbitone, alprazolam, amylobarbitone, aprobarbital, barbital, barbitone, benzphetamine, brallobarbital, bromazepam, brotizolam, buspirone, butalbital, butobarbitone, butorphanol, camazepam, captodiamine, carbromal, carfentanyl, carpipramine, cathine, chloral, chloral betaine, chloral hydrate, chloralose, chlordiazepoxide, chlorhexadol, chlormethiazole edisylate, chlormezanone, cinolazepam, clobazam, potassium clorazepate, clotiazepam, cloxazolam, cyclobarbitone,

delorazepam, dexfenfluramine, diazepam, diethylpropion, difebarbamate, difenoxin, enciprazine, estazolam, ethyl loflazepate, etizolam, febarbamate, fencamfamin, fenfluramine, fenproporex, fluanisone, fludiazepam, flunitraam, flunitrazepam, flurazepam, flutoprazepam, gepirone, glutethimide, halazepam, haloxazolam, hexobarbitone, ibomal, ipsapirone, ketazolam, lorazepam mesylate, lorazepam, lormetazepam, mazindol, mebutamate, medazepam, mefenorex, mephobarbital, meprobamate, metaclozepam, methaqualone, methohexital, methylpentynol, methylphenobarbital, midazolam, milazolam, morphine, nimetazepam, nitrazepam, nordiazepam, oxazepam, oxazolam, paraldehyde, pemoline, pentobarbitone, pentazocine, pentobarbital, phencyclidine, phenobarbital, phendimetrazine, phenmetrazine, phenprobamate, phentermine, phenylacetone, pinazepam, pipradol, prazepam, proxibarbal, quazepam, quinalbarbitone, secobarbital, secbutobarbitone, sibutramine, temazepam, tetrazepam, triazolam, triclofos, zalepan, zaleplon, zolazepam, zolpidem, and zopiclone.

9. The composition of Claim 2, wherein the lipophilic derivative of a drug is a free base or a free acid of the drug.

10. The composition of Claim 2, wherein the lipophilic derivative of a drug is a salt comprising the ionized drug and a lipophilic counter-ion.

11. The composition of Claim 2, wherein the lipophilic derivative of a drug is a complex comprising one or more components selected from the group consisting of drug molecules, metal cations, and lipophilic counter-ions.

12. The composition of Claim 2, wherein the lipophilic derivative of a drug is a complex comprising one or more components selected from the group consisting of drug molecules, metal cations, and cyclodextrin molecules.

13. The composition of Claim 2 wherein the drug is complexed with a metal cation selected from the group consisting of zinc, calcium, magnesium, bismuth and combinations thereof.

14. The composition of Claim 11 wherein the drug is oxycodone,
and

wherein the metal cation is selected from the group consisting of zinc, calcium, magnesium, and bismuth and the lipophilic counter-ion is selected from the group consisting of stearate, oleate, palmitate, laurate and linoleate.

15. The composition of Claim 2, wherein the lipophilic derivative of a drug is a complex comprising the drug and a cyclodextrin.

16. The composition of Claim 2, wherein the lipophilic derivative of a drug is an ester or amide formed between the drug and a fatty acid.

17. The composition of Claim 1, wherein the drug is incorporated into a plurality of individual microparticles comprising a material that is either slowly soluble in water or water insoluble.

18. The composition of Claim 17 wherein the microparticles comprise a wax or wax-like material.

19. The composition of Claim 17 wherein the microparticles comprise a fat or a fatty substance.

20. The composition of Claim 17 wherein the microparticles comprise a material selected from the group consisting of naturally water insoluble proteins, naturally water insoluble polysaccharides, naturally water insoluble lipids and phospholipids, cross-linked water soluble proteins, cross-linked water soluble polysaccharides, cross-linked water soluble cyclodextrins and combinations thereof.

21. The composition of Claim 17 wherein the individual microparticles are coated with one or more independent layers, where at least one of the layers is water insoluble and is degraded by enzymes of the human gastrointestinal tract.

22. The composition of Claim 1 wherein the drug is in the form of individual drug particles coated with one or more independent layers where at least one of the layers is water insoluble and is degraded by enzymes of the human gastrointestinal tract.

23. The composition of Claim 21 wherein at least one of the layers is water-insoluble, organic solvent-insoluble, and degradable by enzymes present in the human gastrointestinal tract.

24. The composition of Claim 21 comprising materials wherein a combination of these materials is not soluble in water, organic solvent, or any combination thereof.

25. The composition of Claim 21 wherein the composition is not completely soluble, and

wherein the drug is not fully released in a single solvent or enzyme solution.

26. The composition of Claim 21 wherein the enzymatically degradable layer(s) comprise a material selected from the group consisting of naturally water insoluble proteins, naturally water insoluble polysaccharides, naturally water insoluble lipids and phospholipids, cross-linked proteins, cross-linked polysaccharides, and combinations thereof.

27. The composition of Claim 1 wherein the drug prone to abuse is co-administered with a drug that has no appreciable abuse potential.

28. The composition of claim 1 formulated for the drug to be immediately released upon oral administration.

29. The composition of claim 1 wherein the drug prone to abuse is oxycodone.

30. A method of manufacturing an abuse-resistant pharmaceutical composition comprising homogeneously dispersing a therapeutically effective amount of a drug prone to abuse, in one or more pharmaceutically acceptable carrier(s), diluent(s), and/or additives, to form an orally administerable abuse-deterrent pharmaceutical composition comprising a therapeutically effective amount of a drug prone to abuse selected from the group of compositions consisting of

(a) a composition comprising a therapeutically effective amount of a lipophilic derivative of a drug prone to abuse, and

(b) a water-insoluble, preferably lipophilic, formulation comprising a therapeutically effective amount of a drug prone to abuse, as defined by any of claims 1-29.

31. The method of claim 30 further comprising formulating the composition into a capsule or tablet.

32. A method of administering an abuse-resistant pharmaceutical composition comprising orally administering to a patient in need thereof an abuse-deterrent pharmaceutical composition comprising a therapeutically effective amount of a drug prone to abuse selected from the group of compositions consisting of

(a) a composition comprising a therapeutically effective amount of a lipophilic derivative of a drug prone to abuse, and

(b) a water-insoluble, preferably lipophilic, formulation comprising a therapeutically effective amount of a drug prone to abuse, as defined by any of claims 1-29.




Figure 1A




Figure 1B




Figure 1C

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/21095

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 9/14, 9/20, 9/24, 9/48,
US CL : 424/458, 459, 468, 472, 475, 490

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
U.S. : 424/458, 459, 468, 472, 475, 490

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
WEST

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X — A	US 6,309,668 B1 (BASTIN et al.) 30 October 2001 (30.10.2001) , see columns 1-2, examples 1-9.	1-9, 17-18, 20-28, 30-32 ----- 10-16, 19, 29

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

29 September 2003 (29.09.2003)

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

Facsimile No. (703)305-3230

Date of mailing of the international search report

06 NOV 2003

Authorized officer

YOTHASNA A VENKAT

Telephone No. 703-308-1235